

**CERTIFICATE OF MAILING BY "EXPRESS MAIL"**

"Express Mail" Mailing Label Number

EL 609 734 209US

Date of Deposit: January 16, 2001

I hereby certify that this paper of fee and the papers indicated as being attached hereto are being deposited with the United States Postal Service "Express mail Post Office to Address" Service under 37 C.F.R. § 1.10 on the date indicated above and are addressed to the Commissioner of Patents and Trademarks, Washington, D.C. 20231.

Patricia Munoz

(Typed or printed name of person mailing)

(Signature of person mailing)

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE  
REQUEST FOR FILING NATIONAL PHASE OF  
PCT APPLICATION UNDER 35 U.S.C. 371 AND 37 CFR 1.494 OR 1.495**

To: Hon. Commissioner of Patents  
Washington, D.C. 20231

TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)

Atty Dkt: PM 099750 /0211500  
M# /Client Ref.

From: Pillsbury Winthrop LLP, IP Group:

Date: January 16, 2001

This is a **REQUEST** for **FILING** a PCT/USA National Phase Application based on:

International Application	2. International Filing Date	3. Earliest Priority Date Claimed
<u>PCT/AU99/00567</u>	<u>14 July 1999</u>	<u>14 July 1998</u>
<u>↑ country code</u>	Day <u>MONTH</u> Year	Day <u>MONTH</u> Year
		(use item 2 if no earlier priority)

Measured from the earliest priority date in item 3, this PCT/USA National Phase Application Request is being filed within:

(a) ☐ 20 months from above item 3 date (b) ☒ 30 months from above item 3 date,

(c) Therefore, the due date (unextendable) is January 16, 2001, since January 14, 2001 was a Sunday and January 15, 2001 was a federal holiday.

5. Title of Invention ANHYDRIDE MODIFIED CANTHARIDIN ANALOGUES USEFUL IN THE TREATMENT OF CANCER

6. Inventor(s) McCLUSKEY, Adam; SAKOFF, Jennette, A.; ACKLAND, Stephen; SIM, Alistair, T.R.

Applicant herewith submits the following under 35 U.S.C. 371 to effect filing:

7. ☒ Please immediately start national examination procedures (35 U.S.C. 371 (f)).
8. ☐ A copy of the International Application as filed (35 U.S.C. 371(c)(2)) is transmitted herewith (file if in English but, if in foreign language, file only if not transmitted to PTO by the International Bureau) including:
- a. ☐ Request;
  - b. ☐ Abstract;
  - c. ☐ pgs. Spec. and Claims;
  - d. ☐ sheet(s) Drawing which are ☐ informal ☐ formal of size ☐ A4 ☐ 11"
9. ☒ A copy of the International Application has been transmitted by the International Bureau.
10. A translation of the International Application into English (35 U.S.C. 371(c)(2))
- a. ☐ is transmitted herewith including: (1) ☐ Request; (2) ☐ Abstract;
  - (3) ☐ pgs. Spec. and Claims;
  - (4) ☐ sheet(s) Drawing which are: ☐ informal. ☐ formal of size ☐ A4 ☐ 11"
  - b. ☒ is not required, as the application was filed in English.
  - c. ☐ is not herewith, but will be filed when required by the forthcoming PTO Missing Requirements Notice per Rule 494(c) if box 4(a) is X'd or Rule 495(c) if box 4(b) is X'd.

RE: USA National Filing of PCT /AU99/00567

d. ☐ Translation verification attached (not required now).

JC07 Rec'd PCT/PTO

16 JAN 2001

11. ☒ **PLEASE AMEND** the specification before its first line by inserting as a separate paragraph:  
 a. ☒ --This application is the national phase of international application PCT/AU99/00567 filed 14 July 1999 which designated the U.S. and that international application ☒ was ☐ was not published under PCT Article 21(2) in English.--  
 b. ☐ --This application also claims the benefit of U.S. Provisional Application No. 60/\_\_\_\_, filed \_\_\_\_.--
12. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3)), i.e., before 18th month from first priority date above in item 3, are transmitted herewith (file only if in English) including:
13. ☒ PCT Article 19 claim amendments (if any) have been transmitted by the International Bureau
14. ☐ Translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)), i.e., of claim amendments made before 18th month, is attached (required by 20th month from the date in item 3 if box 4(a) above is X'd, or 30th month if box 4(b) is X'd, or else amendments will be considered canceled).
15. **A declaration of the inventor** (35 U.S.C. 371(c)(4))  
 a. ☐ Is submitted herewith ☐ Original ☐ Facsimile/Copy  
 b. ☒ Is not herewith, but will be filed when required by the forthcoming PTO Missing Requirements Notice per Rule 494(c) if box 4(a) is X'd or Rule 495(c) if box 4(b) is X'd.
16. **An International Search Report (ISR):**  
 a. Was prepared by ☐ European Patent Office ☐ Japanese Patent Office ☐ Other  
 b. ☒ Has been transmitted by the international Bureau to PTO.  
 c. ☐ Copy herewith (\_\_\_\_ pg(s).) ☐ plus Annex of family members (\_\_\_\_ pg(s).).
17. **International Preliminary Examination Report (IPER):**  
 a. ☒ has been transmitted (if this letter is filed after 28 months from date in item 3) in English by the International Bureau with Annexes (if any) in original language.  
 b. ☐ copy herewith in English.  
 c.1 ☐ IPER Annex(es) in original language ("Annexes" are amendments made to claims/spec/drawings during Examination) including attached amended:  
 c.2 ☐ Specification/claim pages #\_\_\_\_ claims #\_\_\_\_  
 Dwg Sheets #\_\_\_\_  
 d. ☐ Translation of Annex(es) to IPER (required by 30<sup>th</sup> month due date, or else annexed amendments will be considered canceled).
18. **Information Disclosure Statement** including:  
 a. ☐ Attached Form PTO-1449 listing documents  
 b. ☐ Attached copies of documents listed on Form PTO-1449  
 c. ☒ A concise explanation of relevance of ISR references is given in the ISR.
19. ☐ **Assignment** document and Cover Sheet for recording are attached. Please mail the recorded assignment document back to the person whose signature, name and address appear at the end of this letter.
20. ☐ Copy of Power to IA agent.
21. ☐ **Drawings** (complete only if 8d or 10a(4) not completed): \_\_\_\_ sheet(s) per set: ☐ 1 set informal; ☐ Formal of size ☐ A4 ☐ 11"
22. Small Entity Status ☐ is **Not** claimed ☐ is claimed (pre-filing confirmation required)  
 22(a) \_\_\_\_ (No.) Small Entity Statement(s) enclosed (since 9/8/00 Small Entity Statements(s) not essential to make claim)
23. **Priority** is hereby claimed under 35 U.S.C. 119/365 based on the priority claim and the certified copy, both filed in the International Application during the international stage based on the filing in (country) Australia of:
- |     | Application No. | Filing Date  |     | Application No. | Filing Date |
|-----|-----------------|--------------|-----|-----------------|-------------|
| (1) | PP 4665         | 14 July 1998 | (2) |                 |             |
| (3) |                 |              | (4) |                 |             |
| (5) |                 |              | (6) |                 |             |
- a. ☒ See Form PCT/IB/304 sent to US/DO with copy of priority documents. If copy has not been received, please proceed promptly to obtain same from the IB.  
 b. ☐ Copy of Form PCT/IB/304 attached.

09/743841

24. Attached:

JC07 Rec'd PCT/PTO

16 JAN 2001

25. Preliminary Amendment:

25.5 Per Item 17.c2, cancel original pages #\_\_, claims #\_\_, Drawing Sheets #26. **Calculation of the U.S. National Fee (35 U.S.C. 371 (c)(1)) and other fees is as follows:**Based on amended claim(s) per above item(s) ☐ 12, ☐ 14, ☐ 17, ☐ 25, ☐ 25.5 (hilit) ☐

Total Effective Claims	35	minus 20 =	15	x \$18/\$9	=	\$270	966/967
Independent Claims	8	minus 3 =	5	x \$80/\$40	=	\$400	964/965
If any proper (ignore improper) Multiple Dependent claim is present,				add \$270/\$135		+270	968/969

BASIC NATIONAL FEE (37 CFR 1.492(a)(1)-(4)): →→ BASIC FEE REQUIRED, NOW →→→→A. If country code letters in item 1 are not "US", "BR", "BB", "TT", "MX", "IL", "NZ", "IN" or "ZA"

See item 16 re:

1. Search Report was <u>not</u> prepared by EPO or JPO -----	add \$1000/\$500		960/961
2. Search Report was prepared by EPO or JPO -----	add \$860/\$430	+860	970/971

SKIP B, C, D AND E UNLESS country code letters in item 1 are "US", "BR", "BB", "TT", "MX", "IL", "NZ", "IN" or "ZA"

→ <input type="checkbox"/> B. If <u>USPTO</u> did not issue <u>both</u> International Search Report (ISR) <u>and</u> (if box 4(b) above is X'd) the International Examination Report (IPER), -----	add \$970/\$485	+0	960/961
(only) <input type="checkbox"/> C. If <u>USPTO</u> issued ISR but not IPER (or box 4(a) above is X'd), -----	add \$710/\$355	+0	958/959
(one) <input type="checkbox"/> D. If <u>USPTO</u> issued IPER but IPER Sec. V boxes <u>not all</u> 3 YES, -----	add \$690/\$345	+0	956/957
(of) <input type="checkbox"/> E. If international preliminary examination fee was paid to <u>USPTO</u> and Rules 492(a)(4) and 496(b) <u>satisfied</u> (IPER Sec. V <u>all</u> 3 boxes YES for <u>all</u> claims), -----	add \$100/\$50	+0	962/963

27. SUBTOTAL = \$1800

28. If Assignment box 19 above is X'd, add Assignment Recording fee of ----\$40 +0 (581)

29. Attached is a check to cover the ----- TOTAL FEES \$1800

Our Deposit Account No. 03-3975

Our Order No. 099750 C# 0211500 M#

**CHARGE STATEMENT:** The Commissioner is hereby authorized to charge any fee specifically authorized hereafter, or any missing or insufficient fee(s) filed, or asserted to be filed, or which should have been filed herewith or concerning any paper filed hereafter, and which may be required under Rules 16-18 and 492 (missing or insufficient fee only) now or hereafter relative to this application and the resulting Official document under Rule 20, or credit any overpayment, to our Account/Order Nos. shown above for which purpose a duplicate copy of this sheet is attached.

This CHARGE STATEMENT does not authorize charge of the issue fee until/unless an issue fee transmittal form is filedPillsbury Winthrop LLP  
Intellectual Property Group50 Fremont Street  
San Francisco, CA 94105-2228  
Tel: (415) 983-1000

By Atty: Jane K. Babin

Reg. No. 47,224

Sig: Fax: (415) 983-1200  
Tel: (858) 509-4022

Atty/Sec: JRW/JKB/pm

NOTE: File in duplicate with postcard receipts (PAT-103) & attachments.

ANHYDRIDE MODIFIED CANTHARIDIN ANALOGUES USEFUL IN THE TREATMENT OF CANCER**TECHNICAL FIELD**

This invention relates to compounds useful in the treatment of certain forms of cancer; processes for producing these compounds; methods of treatment using these compounds *per se*; methods of treatment using these compounds which methods also increase the sensitivity of cancer cells to other treatments; methods of screening these compounds for anti-cancer activity; and methods of screening these compounds for anti-cancer activity and/or ability to sensitise cancer cells to other methods of treatment. More particularly, the compounds are specific inhibitors of protein phosphatases 1 and 2A.

**BACKGROUND ART**Protein phosphatase inhibitors and the abrogation of cell cycle checkpoints

The regulation of protein phosphatases is integral to the control of many cell processes, including cell growth, transformation, tumour suppression, gene transcription, apoptosis, cellular signal transduction, as neurotransmission, muscle contraction, glycogen synthesis, and T-cell activation. The role of protein phosphatases in many of these processes is often mediated via alterations in the cell cycle. Cell cycle progression is tightly regulated to ensure the integrity of the genome. During cell division it is imperative that each stage of the cell cycle be completed before entry into the next, and this is achieved through a series of checkpoints. The cell cycle can be broken down into four phases, the first gap ( $G_1$ ), is followed by a phase of DNA synthesis (S-phase); this is followed by a second gap ( $G_2$ ) which in turn is followed by mitosis (M) which produces two daughter cells in  $G_1$ . There are two major control points in the cell cycle, one late in

- 2 -

G<sub>1</sub>, and the other at the G<sub>2</sub>/M boundary. Passage through these control points is controlled by a universal protein kinase, cdk1. The kinase activity of cdk1 is dependant on phosphorylation and the association with a regulatory subunit, cyclin B. The periodic association of different cyclins with different cyclin dependent kinases (cdk) has been  
5 shown to drive different phases of the cell cycle; thus cdk4-cyclin D1 drives cells through mid G<sub>1</sub>, cdk2-cyclin E drives cells in late G<sub>1</sub>, cdk2-cyclin A controls entry into S-phase and cdc2-cyclin B drives the G<sub>2</sub>/M transition (O'Connor, 1996, 1997).

Following DNA damage induced by chemotherapy or radiation treatment these checkpoints are responsible for halting cell cycle progression in G<sub>1</sub>, S and/or G<sub>2</sub> phases  
10 (O'Connor, 1996). The cell undergoes a cell cycle arrest so that the damaged DNA can be repaired before entry into S phase or mitosis. The phase at which the cell cycle is halted will depend upon the type of DNA damaging agent used and the point during the cell cycle that the damage was incurred (O'Connor, 1997). The cell cycle is controlled and regulated by an intricate phosphorylation network (Stein et al., 1998). More  
15 particularly, activation of cdk/cyclin complexes requires the phosphorylation of a conserved threonine residue, which are catalysed by CAK kinase, as well as the removal of inhibitory phosphorylations by the phosphatase cdc25. Cdc25 is only active in its phosphorylated form. Therefore, protein phosphatase 2A (PP2A) can inhibit the activation of cdk/cyclin complexes by inhibiting CAK activity and by dephosphorylating  
20 cdc25. The G<sub>1</sub>/S checkpoint is predominantly regulated by the cdk/cyclin D/E complex that mediates its effects by phosphorylating and inactivating the tumour suppressor protein retinoblastoma (pRb). The phosphorylation of pRb prevents it from interacting with the S-phase transcription factor E2F. E2F controls the transcription of proteins

- 3 -

needed for DNA synthesis and entry into S-phase including thymidylate synthase.

Accordingly, the inactivation of pRb by phosphorylation permits entry into the S-phase and vice versa. However, protein phosphatase 1 (PP1) can dephosphorylate pRb and inhibit the cell cycle (Durfee et al., 1993). Thus, PP1 and PP2A are both negative  
5 regulators of the cell cycle. Inhibition of PP1 and PP2A would abrogate these checkpoints and prematurely force cells through the cell cycle.

Serine/threonine phosphatases, which are responsible for protein dephosphorylation, comprise a unique class of enzymes consisting of four primary subclasses based on their differences in substrate specificity and environmental  
10 requirements. Of the serine/threonine phosphatases, protein phosphatases 1 and 2A (PP1 and PP2A, respectively) share sequence identity between both enzyme subunits (50% for residues 23-292; 43% overall), are present in all eukaryotic cells and are together responsible for 90% of all cellular dephosphorylation. Knowledge of structure and subsequent correlation of binding function for both PP1 and PP2A would therefore  
15 provide a vital link toward understanding the biochemical role of these enzymes. A goal of the medicinal chemist is the development of potent and selective inhibitors of these protein phosphatases.

The natural toxins, okadaic acid, calyculin A, microcystin-LR and tautomycin are representative of a structurally diverse group of compounds that are all potent protein  
20 phosphatase 1 (PP1) and 2A (PP2A) inhibitors. Okadaic acid is more specific for PP2A ( $IC_{50}$  1nM) than PP1 ( $IC_{50}$  60nM), while calyculin is slightly more specific for PP1 ( $IC_{50}$  0.5-1.0nM) than PP2A ( $IC_{50}$  2nM). All of these phosphatase inhibitors are known to abrogate cell cycle checkpoints, particularly the  $G_2$  checkpoint of the cell cycle and

- 4 -

induce cellular mitoses (Yamashita et al., 1990). Abrogation of the G<sub>2</sub> checkpoint means that the cell does not have the capacity to detect DNA damage or malformation of the genome prior to entry into mitosis. Therefore, cells which have a deficient G<sub>2</sub> checkpoint are unstable, and incapable of detecting DNA damage, initiating G<sub>2</sub> arrest, or undergoing DNA repair. Such cells enter the mitotic stage of the cell cycle prematurely with malformed spindles. The abrogation is of the G<sub>2</sub> checkpoint in the cell cycle by okadaic acid is mediated via the activation of cdc2/H1 kinase, the major mitotic inducer, and results in a premature mitotic state (Yamashita et al., 1990). Although okadaic acid is known as a tumour promoter, in some cell types, it has been shown to revert the phenotype of oncogene-transformed cells to that of normal cells, and to inhibit neoplastic transformation of fibroblasts (Schonthal, 1991).

Furthermore, okadaic acid has been shown to selectively enhance the cytotoxicity of vinblastine and the formation of apoptotic cells, in HL60 cells which are p53 nul (Kawamura, 1996). Interestingly, calyculin enhances irradiation killing in fibroblast cells at doses that are non toxic when given as a single treatment. (Nakamura and Antoku, 1994). Data also shows that okadaic acid can abrogate the G<sub>1</sub>/S checkpoint of the cell cycle. In this context, okadaic acid has been shown to override the S-phase checkpoint and accelerate progression of G<sub>2</sub>-phase to induce premature mitosis (Gosh et al., 1996). In addition, okadaic acid has been shown to significantly increase the fraction of quiescent cells entering the S-phase via modifications in the phosphorylation state of pRb (Lazzereschi et al., 1997). Other studies have shown that the hyperphosphorylation state of pRb forces cells prematurely into S-phase and pRb can be kept in a phosphorylated state via protein phosphate inhibition (Herwig and Strauss, 1997). Cells

- 5 -

lacking functional pRb show increased apoptosis and cytotoxicity following 5-fluorouracil and methotrexate treatment (Herwig and Strauss, 1997). We propose that cell death would be substantially enhanced in cells forced to enter the S-phase prematurely (via G<sub>1</sub> checkpoint abrogation) and which were lacking key S-phase components such as dTMP (via TS inhibition).

The okadaic acids class of compounds, with the exceptions of okadaic acid, cantharidin (Honaken) and thyriferyl 23-acetate (Matszawa et. al) (being PP2A selective) exhibit poor selectivity. Furthermore, the concentration of PP1 and PP2A inside cells is such that high concentrations of these inhibitors are required to generate a response *in vivo* resulting in the loss of effectiveness of any *in vitro* selectivity (Wang).

Cantharidin (exo,exo-2,3-dimethyl-7-oxobicyclo[2.2.1]heptane-2,3-dicarboxylic acid anhydride), is a major component of the Chinese blister beetles: *Mylabris phalerata* or *M. cichorii* (Yang; Cavill et. al). The dried body of these beetles has been used by the Chinese as a natural remedy for the past 2000 years. Although Western medicine decreed cantharidin to be too toxic in the early 1900's (Goldfarb et. al) its purported aphrodisiac qualities (the active ingredient of "Spanish Fly"), and its widespread occurrence in cattle feed still results in numerous human and livestock poisonings (Schmitz).

Li and Casida, and previous work in this laboratory (McCluskey et. al) (and more recently Pombo-Villar, Sodeoka) has assisted in the delineation of certain features crucial for inhibition of PP2A by cantharidin analogues (Figure 1). However the corresponding picture for PP1 is not so clear, the majority of data refers to possible



interactions with the known crystal structures, and in some cases the inhibition values for PP1 are not reported.

#### Involvement of Tumour Suppressor Gene p53

The most commonly mutated gene in human cancers is the tumour suppressor gene p53, which is abnormally expressed in more than 50% of tumours. The development of chemotherapeutic agents which selectively target cancer cells with mutant p53 is certainly desirable, for two main reasons. Firstly, cells that have an abnormal p53 status are inherently resistant to conventional chemotherapy and produce the more common, and more aggressive tumours such as colon carcinoma and non small cell lung cancer. Secondly, a chemotherapy regime that targeted only those cells with a mutant p53 phenotype would potentially produce fewer side effects since only the cancer cells would be killed and not the p53 proficient normal healthy cells.

### **DISCLOSURE OF THE INVENTION**

In relation to the discussion above, the present inventors believed that the replacement of the ether O atom of the anhydride with N or S (as N-H and N-R, where R = alkyl or aryl) would allow them to probe the H-bonding requirements of this region of cantharidin analogues. Previous studies in their laboratory had shown limited tolerance for modification of the 7-oxa position. An ability to modify these heteroatoms is crucial to the development of selective inhibitors based on this simple skeleton.

There is not, at present, an inhibitor with either absolute specificity or high enough selectivity which renders the inhibitor effectively specific in vivo.


It has surprisingly been found that anhydride modified cantharidin analogues, which are the subject of this invention, may possess one or more of the properties of

- 7 -


being potent, selective, oxidatively stable, and cell permeable inhibitors of protein phosphatases 1 and 2A.

Therefore, according to the first aspect of this invention there are provided cell permeable inhibitors of protein phosphatases 1 and 2A, said inhibitors being anhydride  
5 modified cantharidin analogues.

According to a particular embodiment of the first aspect of this invention there are provided compounds of the formula:



wherein  $R_1$  and  $R_2$  are H, aryl or alkyl; X is O, N or S; Y is O, S, SR, NH, NR,  $CH_2OH$ ,  
10  $CH_2OR$ ; R is alkyl or aryl; A and B are H or  $CH_3$ ; W and Z are  $CHOH$  or  $C=O$  and  $R_1$   
and  $R_2$  can cyclise to form a ring as follows:



wherein  $R_3$  and  $R_4$  are H, aryl or alkyl.

The aryl group may suitably be phenyl or naphthyl for example, and may be  
15 attached via a carbon spacer of between 6 and 10 carbon atoms. The alkyl group may  
suitably be  $C_1$ - $C_{10}$ .

- 8 -

According to the second aspect of this invention there is provided a process for producing anhydride modified cantharidin analogues. The process may include the steps of:

dissolving a diene in a suitable solvent and adding to the resultant solution an  
5 ene.

According to a third aspect of the invention there is provided a process for producing anhydride modified cantharidin analogues, involving the step of reacting a diene with an ene.

The process may further involve hydrogenation of the adduct of the diene and  
10 ene and/or optionally, ring opening of the adduct.

Generally, the reaction conditions for the production of the anhydride modified cantharidin analogues are dependent on the aromaticity of the starting diene. Suitable reaction conditions are exemplified below.

According to a fourth aspect of this invention there is provided a method of  
15 treating a cancer which method comprises administering to a patient in need of such treatment, an effective amount of an anhydride modified cantharidin analogue of the first aspect of this invention, together with a pharmaceutically acceptable carrier, diluent and/or excipient.

The method may be carried out in conjunction with one or more further  
20 treatments for treating the cancer.

According to a fifth aspect of this invention there is provided a method of sensitising cancer cells to at least one method of treating cancer, which method of sensitising comprises administering to a patient in need of such treatment, an effective

amount of an anhydride modified cantharidin analogue of the first aspect of this invention, together with a pharmaceutically acceptable carrier, diluent and/or excipient.

According to a sixth aspect of the invention there is provided a method of treating cancer which method comprises:

- 5 administering to a patient in need of such treatment, an effective amount of an anhydride modified cantharidin analogue to sensitise cancer cells of the patient to one or more cancer treatments; and utilising the one or more cancer treatments.

According to a seventh aspect of this invention there is provided a method of screening a compound for anti-cancer activity.

- 10 According to an eighth aspect of this invention there is provided a method of screening compounds for use in the fourth aspect of this invention, said method comprising screening for anti-cancer activity; and screening for ability to abrogate either the G<sub>1</sub> or the G<sub>2</sub> checkpoint of the cancer cell cycle. The method may also comprise screening for the ability of said compounds to sensitise cancer cells to one or more  
15 cancer treatments.

The one or more cancer treatments mentioned above may be selected from treatments involving cisplatin, irradiation, taxanes and antimetabolites.

The invention will hereinafter be described with reference to Examples and the accompanying figures.

20 Brief Description of the Figures

Figure 1 is a schematic representation of the structure activity data generated for inhibition by PP2A by cantharidin analogues;

Figure 2: New cantharidin analogues.

Figure 3: Cytotoxicity of cantharidin and the new cantharidin analogues.

Figure 4: Cell cycle analysis 12h following exposure to cantharidin, MK-2 or MK-4.

Figure 5: Cell cycle analysis 18h after 6Gy of radiation and 12h after exposure to cantharidin, MK-2 or MK-4.

5        Figure 6 (a-c): Combination index versus fraction affected: HCT116 colon cells in simultaneous combination with cisplatin and MK-4.

Figure 7 (a-b): Combination index versus fraction affected: HT29 colon cells in simultaneous combination with cisplatin and MK-4.

Figure 8 (a-c): Combination index versus fraction affected: HCT116 colon cells  
10    in simultaneous combination with taxotere and MK-4

Figure 9 (a-c): Combination index versus fraction affected: HT29 colon cells in simultaneous combination with taxotere and MK-4.

#### Best and other Modes for Carrying Out the Invention

As mentioned above, the reaction conditions for producing anhydride modified  
15    cantharidin analogues encompassed by the present invention generally depend on the aromaticity of the starting diene. This is illustrated by a description of examples of the methods wherein the starting materials are furan (Method 1 below); thiophene (Method 2 below); and pyrrole (Method 3 below).

#### Method 1: Furan as the starting diene

20        A solution of furan (5 equivalents) is dissolved in a suitable solvent (about 5 times the volume of furan, the solvent can be for example, ether (for room temperature reactions); or benzene or xylene (the latter two for reactions at 80 and 130°C respectively). To this solution is added one equivalent of the ene. The reaction is then heated (or stirred at room temperature), typically for 24 hours (2 days in the case of the  
25    room temperature reaction). Upon cooling (or standing at room temperature) a precipitate forms and is collected by vacuum filtration. The adduct is then purified by

- 11 -

recrystallisation from for example, chloroform or ethanol. In the case of the furan + maleic anhydride compound care is exercised to minimise heating as this causes a retro-Diels-Alder reaction yielding only the starting materials.

Method 2: Thiophene as the starting diene

5 Thiophene (1.016g, 0.012 mol) and maleic anhydride (0.558.0.006 mol) are mixed at room temperature in 2.5 mL of distilled dichloromethane. This mixture is then placed inside a high pressure reactor. They are compressed to a pressure of 17kbar at 40°C for a period of 71 hours, after which the pressure is released and the product purified by chromatography.

10 Method 3: Pyrrole as the starting diene

To  $[\text{Os}(\text{NH}_3)_5\text{OsO}_2\text{CF}_3] (\text{CF}_3\text{SO}_3)_2$ , (0.3511 g, 0.4 mmol) and activated magnesium (0.1511 g), pyrrole (0.45 mL, 0.6 mmol), DME (1 mL) and DMAc (0.3 mL) are added in that order. The mixture is stirred for 1 hour, the temperature gradually rising to 40°C and then dropping. The brown slurry is filtered through a thin pad of  
15 celite, and the cake washed with DME in small portions (4 x 2 mL). The filtrate is added to dichloromethane (15 mL). Vigorous stirring results in the formation of yellow coloured precipitate which is collected by vacuum filtration, followed by an ether wash (2 x 2.5 mL). The product is dried under a stream of nitrogen yielding a yellow-tan solid (0.343g, 84%). To this pyrrole complex is added maleimide (0.05g, 0.515 mmol)  
20 (or any other "ene", eg maleic anhydride, dimethyl maleate, etc) in acetonitrile. The mixture is allowed to stir at room temperature for 60 min. after which the solvent is removed by vacuum, yielding the exo isomer (0.359g, 64%). The crude material is purified by ion-exchange column (Sephadex-CM C-25, 2 x 10 cm), using NaCl as the

- 12 -

mobile phase. The complexes are precipitated by the addition of a saturated sodium tetraphenylborate solution.

The types of cancer which are amenable to treatment by these compounds include those types of cancer which are inherently resistant to conventional chemotherapy. Typically, these types of cancer are represented by the more common and more aggressive tumour types such as, but not limited to, colon cancer and non small-cell lung cancer.

The compounds of this invention are suitably administered intravenously, although other modes of administration are possible. Pharmaceutically acceptable diluents, adjuvants, carriers and/or excipients may be used in conjunction with the compounds of this invention.

Suitable such pharmaceutically acceptable substances are those within the knowledge of the skilled person and include compounds, materials and compositions deemed appropriate.

Actual dosage levels of the compounds of the invention may be varied so as to obtain an amount of the active ingredient which is effective to achieve the desired response for a particular patient, composition and mode of administration.

The dosage level can be readily determined by the physician in accordance with conventional practices and will depend upon a variety of factors including the activity of the particular compound of the invention to be administered, the route of administration, the time of administration, the rate of excretion of the particular compound employed, the age, sex, weight, condition, general health and prior medical history of the patient being treated, and like factors well known in the medical arts.

The compounds of this invention may also sensitise cancer cells to other methods of treatment. For example, typically these methods include irradiation and treatment with platinum anti-cancer agents, for example cisplatin.

In addition, sensitisation may also be brought about by, for example the use of  
5 the plant alkaloids vinblastine and vincristine, both of which interfere with tubulin and the formation of the mitotic spindle, as well as taxanes and antimetabolites, including 5-fluorouracil, methotrexate and antifolates.

In particular, the compounds of this invention sensitise those cells with deficient p53 activity.

10 When screening for anti-cancer activity as contemplated by the invention, various cancer cell lines may be chosen. These are typically both haematopoietic and solid tumour cell lines with varying p53 status and include: L1210 (murine leukaemia, p53 wildtype), HL60 (human leukaemia, p53 nul), A2780 (human ovarian carcinoma, p53 wildtype), ADDP (cisplatin resistant A2780 cells, p53 mutant), SW480 (human  
15 colon carcinoma, p53 mutant), WiDr (human colon carcinoma, p53 mutant), HT29 (human colon carcinoma, p53 mutant), HCT116 (human colon carcinoma, p53 wildtype) and 143B (human osteosarcoma, p53 mutant).

In addition to the methods for screening for anti-cancer activity, the following procedures may be suitably used in the remainder of the screening process. For  
20 example, when screening for the ability to abrogate the G<sub>1</sub> and/or the G<sub>2</sub> checkpoint of the cancer cell cycle, the following are suitably used:



Cell cycle method

The cells are fixed in 70% ethanol and stored at - 20°C until analysis is performed (1-2 weeks). After fixing, the cells are pelleted and incubated in PBS containing propidium iodide (40mg/ml) and RNase A (200 mg/ml) for at least 30 min at room temperature. The samples ( $2 \times 10^4$  events) are analysed using a Becton Dickson FACSscan, fluorescence is collected in fluorescence detector 2 (FL2), filter 575/30 nm band pass. Cell cycle distribution is assessed using Cell Quest software (Becton Dickson).

Those protein phosphatase inhibitors which show abrogation of either the G<sub>1</sub> or G<sub>2</sub> checkpoint will then be exploited in combination studies with either radiation exposure or chemotherapy drugs incubation. The MTT (3-[4,5-dimethylthiazol-2-yl]2,5-diphenyl-tetrazolium bromide) assay is used to determine whether a synergistic, antagonistic or additive effect is induced. The Median Effect method is adopted to mathematically determine the optimal combination index of the treatments chosen (Chou and Talalay, 1984). This method has been extensively used to investigate the cytotoxicity of various drug combinations including cisplatin and D1694 (Ackland et al 1996; 1998). A combination index value less than 1 indicates synergism, a value equal to 1 indicates additivity and a value greater than one indicates antagonism.

Cytotoxicity assay

When screening for the ability to sensitise cancer cells to conventional chemotherapy and irradiation, the following methods are suitably used:

Cells in a subconfluent phase are transferred to 96- well microtitre plates. L1210 cells are plated at a density of 1000 cells/well in 100µl medium, while all other cell

lines are plated at a density of 2000-25000 cells/well. The cells are left for 24h prior to treatment to ensure exponential growth has been achieved, 24h after plating (day 0), 100µl of phosphatase inhibitor is added to each well, control wells received 100µl of medium only. Drug exposure time is 72h (day 3). The effect of phosphatase inhibition is tested in triplicate over a concentration range of  $1 \times 10^{-3} \text{M}$  -  $1 \times 10^{-8} \text{M}$ . Growth inhibitory effects are evaluated using the MTT assay and absorbance read at 540 nm. The  $\text{IC}_{50}$  is the drug concentration at which cell growth is 50% inhibited based on the difference of optical density on day 0 and day 3 of drug exposure. Cytotoxicity is evaluated using a spectrophotometric assay which determines the percentage of cell growth following exposure of the cells to various concentrations of the phosphatase inhibitors for a period of 72 hours. The subsequent dose response curve is used to calculate  $\text{IC}_{50}$  values (the drug concentration at which cell growth is 50% inhibited).

Most drug discovery has focused on the development of new single agents. However, in light of the success of combination chemotherapy it is increasingly apparent that successful anticancer treatment of the future will be based upon the discovery of agents which are synergistic in their action. In view of this, the cytotoxicity of phosphatase inhibitors in combination with either radiation, cisplatin, taxanes, antimetabolites or plant alkaloids is examined. As indicated above, calyculin which by itself is not cytotoxic, enhances irradiation induced cell death. Similarly abrogation of the  $\text{G}_2$  checkpoint by either, caffeine or UCN-01, also enhances the cytotoxicity of  $\gamma$  irradiation in cells with mutant p53 (CA46 and HT-29 cells) (Powell et al., 1995; Russell et al., 1995; Wang et al., 1996). DNA damage induced by irradiation causes both a  $\text{G}_1$  and  $\text{G}_2$  cell cycle arrest. In p53 mutant cells, the  $\text{G}_1$  checkpoint is absent.

- 16 -

However, following irradiation the cells will still arrest in the G<sub>2</sub> phase, and potentially repair the damage. P53 mutant cells are generally more resistant to conventional chemotherapy and produce more aggressive tumours. Therefore, in p53 deficient cells, DNA damage that is not detected by the G<sub>1</sub> checkpoint will be picked up by the G<sub>2</sub> checkpoint. If the cells are deficient in both of these checkpoints then it is believed that the cells will be unable to initiate repair mechanisms and will be more unstable and increasingly susceptible to cell death induced by DNA damage.


Cisplatin is another commonly used anticancer treatment which binds to DNA and produces DNA crosslinks and strand breaks. Cisplatin is particularly useful in the treatment of testicular carcinoma, small cell carcinoma of the lung, bladder cancer, and ovarian cancer. Repair of cisplatin induced DNA damage is mediated via nucleotide excision repair which is coordinated by p53 activation of Gadd45 (Smith et al., 1994). In this context, it has been suggested that cells that are p53 mutant are more sensitive to cisplatin treatment (Hawkins et al., 1996). A number of researchers have investigated this proposal in p53 mutant cell lines and in p53 mutant tumours, with mixed results. While it is apparent that cisplatin is more cytotoxic in cells lines that are deficient in p53 (induced via papillomavirus) compared to the p53 proficient cells (Hawkins et al., 1996), it is harder to test this hypothesis in tumours and in cisplatin resistant cells as they may have several undefined mutations in their genome which would confound such studies (Herod et al., 1996). Nevertheless, the G<sub>2</sub> abrogator UCN-01 (7-hydroxystaurosporine, a protein kinase inhibitor) has been shown to markedly enhanced the cell-killing activity of cisplatin in MCF-7 cells defective for p53 function (Wang et al., 1996).

The development of chemotherapeutic agents which selectively target p53 mutant cells is desirable since 50% of tumours have either a mutated or deleted p53 gene. Many of these p53 deficient cells and tumours are inherently resistant to conventional chemotherapy and represent the common more aggressive tumour types such as colon cancer, and non-small cell lung cancer. Thymidylate synthase (TS) inhibitors are another class of commonly used anticancer agents. TS catalyses a critical step in the pathway of DNA synthesis by converting dUMP to dTMP by methylation using the co-substrate N5,N10-methylene tetrahydrofolate (CH<sub>2</sub>-THF) as a methyl donor. This step is the only *de novo* source of dTMP, which is subsequently metabolised to dTTP exclusively for incorporation into DNA during synthesis and repair (Jackman & Calvert, 1995). Thus, TS is a key regulatory enzyme during the S-phase of the cell cycle. Lack of dTTP results in DNA damage and ultimately cell death, but the process(es) by which cell death occurs is not clear. TS inhibitors such as fluorouracil, raltitrexed, and LY231514 play a pivotal role in anticancer treatment and are often the first line treatment of many cancers (Peters & Ackland, 1996). We propose that the TS inhibitor Thymitaq (Zarix, Ltd) be used in combination with cantharidin analogues. Thymitaq is a direct and specific TS inhibitor which does not require active transport into the cell nor does it require intracellular activation for its action.


The following examples are not to be construed as limiting on the scope of the invention as indicated above.

Example 1Chemistry


Anhydride modified cantharidin analogues were synthesised by a variety of modified literature procedures, as set out in schemes 1 and 2. These modifications are embodied in the three methods, which depend on the aromaticity of the starting dienes, set out above. The dimethyl ester (3), which was prepared by the application of high pressure, 17kbar, 40°C, 61 hours, as shown in scheme 3.



- 10 **Scheme 1.** a. Furan:maleic anhydride (5:1). diethylether, 2d. RT, 96%; b. H<sub>2</sub> / 10% Pd-C/ EtOH; c. p-TosOH, MeOH, chromatography; d. H<sub>2</sub> / 10% Pd-C/ Acetone; e. NaBH<sub>4</sub> then HCl.



**Scheme 2. Reagents and Conditions:** f. Furan: maleimide (5:1), diethyl ether, 7d, in dark, 75%, exo product; g. Furan: Maleimide (5:1), diethylether, sealed tube 12h, 90°C, 66%,endo product.



5 **Scheme 3. Reagents and Conditions:** h.

Furan:dimethylmaleate (2:1), CH<sub>2</sub>Cl<sub>2</sub> ,17 Kbar, 40°C, 61 h, 56%.

### Example 2


Development of potent, selective, oxidatively stable, and cell permeable inhibitors of protein Phosphatases 1 and 2A.

10 Crude natural product extracts have yielded isopalinurin and a series of cantharidin analogues have been synthesised. In this context, the present inventors have developed the simple cantharidin analogue which is PP1 selective (IC<sub>50</sub> = 50mM, with 0% inhibition of PP2A at concentrations  $\geq 1000\text{mM}$ ) representing the first small molecule to exhibit selectivity for PP1. Results have indicated that a series of simple  
15 synthetic modification of the cantharidin skeleton also allows the synthesis of a PP2A selective compound (see Figure 1).

The present inventors have previously demonstrated that a facile ring opening of an anhydride is crucial to inhibition of PP2A. This is not possible with c (previous studies with the 7-0, and this analogue indicated considerable hydrolytic stability of the  
20 maleimide link). It is also interesting to note that endothal thioanhydride is three fold more potent than cantharidin, with the S atom being an important factor. It is thus

envisaged that the 7-S group presents itself to the active sites metals and the N-H of the maleimide occupies the hydrogen bond cavity normally reserved for the 7-O substituent cantharidin.

Structure of cantharidin and selective analogues




- (a) Shows structure of cantharidin;  
(b) Shows PP1 selective analogue; and  
(c) Shows PP2A selective analogue. In the case of panel (c)  $IC_{50} \sim 25\text{mM}$ .

On the basis of these results and previous experience in our laboratory (synthesis and molecular modelling of cantharidin inhibitors at PP1 and PP2A), we have designed a series of analogues which are more active and selective, whilst retaining the desirable properties of stability and cell permeability.

The synthetic pathways to these analogues are shown in schemes 1-3. Each scheme allows for modification of the basic skeleton, and in some cases the insertion of beneficial feature that were present in the more complex natural toxin(s) (eg okadaic acid, calyculin, microcystin, etc). The inclusion of these features is designed to provide enhanced selectivity and potency.

- 21 -

Example 3Synthetic development of a series of PP1 and PP2A analogues of cantharidin.


- (i) Diels-Alder addition (maleic anhydride) and subsequent manipulations of X;
- 5 (ii) Diels-Alder addition (substituted maleic anhydrides), introduction and manipulation of Z (Z = hydrophobic tail; eg long chain nitrile: cf Calyculin A, long chain terminating in a spiro acetal: cf Tautomycin. Okadaic acid; long chain terminating in an aromatic ring: cf Adda in Microcystin-LR; (iii) stereospecific ring opening of the anhydride allowing further manipulations of the newly released functional groups (see scheme 2).

10 In this instance we have developed synthetic protocols in our laboratory that allow the facile assembly of these analogues. Biological evaluation and molecular modelling of the most active molecules will allow compounds to be evaluated.


Additional modification to the basic structure can be obtained as exemplified below.



- 22 -

Example 4

A specific example of one class of cantharidin analogue that shows promise as a  
 5 selective inhibitor of protein phosphatases 1 and 2A.

Example 5Stereospecific route towards 7-azabicyclo [2.2.1] heptanes

We have shown that the introduction of the bridgehead nitrogen improves the  
 10 potency, selectivity and stability of similar analogues. the above pathway has been  
 developed to further improve the bio-activity of these analogues. The synthetic routes  
 alluded to herein may allow the rapid assembly of the target molecules.

Those agents which meet the requirements of being stable, specific, potent, and  
 membrane permeable protein phosphatase inhibitors are screened for their anti-cancer  
 15 activity.

### Example 6

#### Biochemistry

All synthesised compounds were tested for their ability to inhibit protein phosphatases 1 and 2A. Initial investigations were carried out at 100 mM. Promising analogues were then assayed in triplicate for estimation of IC<sub>50</sub> values.

Protein phosphatase 1 and 2A were partially purified from chicken skeletal muscle essentially as described by Cohen. Protein phosphatase activity was measured at 37°C in 50 mM Tris-HCl buffer (pH 7.4), 0.1 mM EDTA, 5 mM caffeine, 0.1% 2-mercaptoethanol and 1 mg/ml bovine serum albumin using 30 mg [<sup>32</sup>P]-phosphorylase as substrate. The total assay volume was 30 ml. The assay conditions were restricted to 20% dephosphorylation to ensure linearity and inhibition of protein phosphatase activity was determined by including cantharidin or its analogues at the required concentrations in the reaction buffer. Reactions were terminated by the addition of 0.1 ml ice cold 20% trichloroacetic acid. Precipitated protein was pelleted by centrifugation and the radioactivity in the supernatant measured by liquid scintillation counting. Data is expressed as the percentage inhibition with respect to a control (absence of a competing compound) incubation.

### Example 7

#### Screening various PP1 and PP2A inhibitors for anti-cancer activity

(a) Cytotoxicity of protein phosphatase inhibition:

Those PP1 and PP2A inhibitors which fulfil the requirements detailed above were tested in various cancer cell lines. The cell lines chosen for study included both haematopoietic and solid tumour cell lines with varying p53 status and include:

- 24 -

L1210 (murine leukaemia, p53 wildtype),  
 HL60 (human leukaemia, p53 nul),  
 A2780 (human ovarian carcinoma, p53 wildtype),  
 ADDP (cisplatin resistant A2780 cells, p53 mutant),  
 5 SW480 (human colon carcinoma, p53 mutant),  
 WiDr (human colon carcinoma, p53 mutant).  
 HT29 (human colon carcinoma, p53 mutant)  
 HCT116 (human colon carcinoma, p53 wildtype)  
 143B (human osteosarcoma, p53 mutant)

10 Anti-cancer screening of the protein phosphatase inhibitors is assessed using the  
 MTT assay. This assay determines cell viability by the ability of mitochondrial  
 dehydrogenase to produce formazan crystals from 3-(4,5-dimethylthiazol-2-yl) -2, 5-  
 diphenyltetrazolium bromide. The viable cell number/well is directly proportional to the  
 production of formazan, which following solubilization, can be measured  
 15 spectrophotometrically (540nm). This technique is also used by the National Cancer  
 Institute to screen for new anticancer agents.

As described herein a number of cantharidin analogues have been synthesised  
 and tested for their anticancer activity in nine cancer cell lines using the MTT assay after  
 72 h exposure. These new analogues are shown in Figure 2 and have been designated  
 20 MK-1 through to MK-9. The cytotoxicity ( $IC_{50}$ ) of these cantharidin analogues is shown  
 in Table 1 and Figure 3. In summary, the MK-1 analogue did not show any significant  
 cytotoxicity in any of the cell lines tested ( $IC_{50} > 1000\mu M$ ). Only marginal cytotoxicity  
 across all cell lines tested was observed for MK-3 ( $IC_{50}$  247 to  $> 1000\mu M$ ), MK-7 ( $IC_{50}$

Table 1

IC<sub>50</sub> values of tumour cell lines after 72 h continuous exposure to cantharidin and cantharidin analogues.

Tumour type	Cell line	p53 status	IC <sub>50</sub> (mean ± SE) after 72h continuous exposure (μM)								
			Cantharidin	MK-1	MK-2	MK-3	MK-4	MK-5	MK-7	MK-8	MK-9
Murine Leukaemia	L1210	wt	18 ±	>1000	185 ± 51	647 ± 132	680 ± 97	>1000	367 ± 37	337 ± 19	192 ± 56
Human Leukaemia	HL60	nul	13 ±	>1000	177 ± 3	247 ± 55	393 ± 103	323 ± 13	293 ± 7	297 ± 3	133 ± 9
Human Ovarian	A2780	wt	±	>1000	157 ± 9	317 ± 17	333 ± 55	567 ± 109	357 ± 102	313 ± 61	187 ± 9
Human Ovarian	ADDP	mt	12 ± 0.8	>1000	183 ± 17	>1000	275 ± 56	260 ± 40	210 ± 18	208 ± 19	233 ± 23
Human Osteosarcoma	143B	mt	10.2 ± 1.2	>1000	248 ± 29	665 ± 225	450 ± 50	>1000	327 ± 67	385 ± 43	223 ± 44
Human Colon	HCT116	wt	12 ±	>1000	160 ± 10	>1000	78 ± 7	143 ± 23	180 ± 20	173 ± 22	107 ± 12
Human Colon	HT29	mt	6.4 ± 0.7	>1000	183 ± 20	530 ± 112	14 ± 0.3	28 ± 1	297 ± 58	373 ± 54	205 ± 13
Human Colon	WiDr	mt	6.1 ± 0.5	>1000	198 ± 53	620 ± 31	15 ± 3	31 ± 10	320 ± 20	367 ± 44	190 ± 35
Human Colon	SW480	mt	17.5 ±	>1000	155 ± 9	444 ± 27	88 ± 5	247 ± 14	333 ± 22	353 ± 20	147 ± 14

wt = wildtype, mt = mutant.

- 26 -

180-367 $\mu$ M) and MK-8 (IC<sub>50</sub> 173-385 $\mu$ M). Greater cytotoxicity was observed with MK-2 (IC<sub>50</sub> 157-248 $\mu$ M) and MK-9 (IC<sub>50</sub> 107-233 $\mu$ M) which was also consistent across the nine cell lines. The greatest cytotoxicity was observed with the MK-4 and MK-5 analogues, however, the magnitude of this response was cell line dependent. In this context, MK-4 and MK-5 were selectively more cytotoxic in the human colon cancer cell lines (IC<sub>50</sub> 14-88 $\mu$ M; 28-247 $\mu$ M) compared with leukaemia (IC<sub>50</sub> 393-680 $\mu$ M; 323->1000 $\mu$ M) ovarian (IC<sub>50</sub> 275-333 $\mu$ M; 260-567 $\mu$ M), and osteosarcoma (IC<sub>50</sub> 450 $\mu$ M; >1000 $\mu$ M) cells respectively.

(b) Abrogation of cell cycle checkpoints:

The ability of the protein phosphatase inhibitors to abrogate the G<sub>1</sub> or G<sub>2</sub> checkpoint of the cell cycle may be determined by cell cycle analysis using flow cytometry. Briefly, asynchronous cell cultures are harvested 18h after 6Gy irradiation and/or 12h incubation with the protein phosphatase inhibitor. Depending upon the p53 status of the cell line, radiation treatment alone will induce arrest in either G<sub>1</sub> and/or G<sub>2</sub> phase of the cell cycle.

Data shown in Table 2 and Figure 4 show the cell cycle response of L1210, HL60, HT29 and HCT116 cells to cantharidin and the new cantharidin analogues MK-2 and MK-4 after 12h exposure. In summary, cantharidin and MK-2 produced a similar response and induced G<sub>2</sub> arrest in all four cell lines tested. MK-4 also induced G<sub>2</sub> arrest but only in L1210, HL60 and HCT116 cells. In HT29 cells, MK-2 induced G<sub>1</sub> cell cycle arrest. The magnitude of the cell cycle arrest induced by these drugs directly correlated with their cytotoxicity in the respective cell lines. The ability of the parent compound cantharidin to inhibit cell growth is also shown (IC<sub>50</sub> 6.1-18 $\mu$ M). The cytotoxicity of the

Table 2

## Cell Cycle Analysis

Cell Cycle Distribution (percentage of total) of tumour cell lines 12h after cantharidin or cantharidin analogue treatment.

Method : Flow Cytometry of Propidium Iodide stained cells.

Agent	L1210 cells					HL60 cells					HCT116 cells					HT29 cells				
	$\mu\text{M}$	sub G <sub>1</sub>	G <sub>1</sub>	S	G <sub>2</sub> +M	sub G <sub>1</sub>	G <sub>1</sub>	S	G <sub>2</sub> +M		sub G <sub>1</sub>	G <sub>1</sub>	S	G <sub>2</sub> +M		sub G <sub>1</sub>	G <sub>1</sub>	S	G <sub>2</sub> +M	
Cantharidin	0	0.5	47.4	34.3	19.4	1.9	45.5	25.8	28.2		6.5	43.3	14.6	36.4		11.1	45.3	8.0	36.0	
	1	0.5	45.8	33.7	21.6	1.5	44.0	26.1	29.7		2.2	39.9	17.2	41.9		9.0	46.2	7.8	37.4	
	5	0.6	46.5	32.6	21.9	1.7	41.4	27.7	30.6		2.9	39.9	16.8	41.8		4.0	47.4	9.3	39.8	
	10	0.5	49.1	33.0	18.9	1.7	41.8	27.5	30.4 G <sub>2</sub> arrest		6.2	38.0	14.9	42.0		2.8	42.7	14.6	40.3 G <sub>2</sub> arrest	
	50	1.9	22.0	27.8	50.6 G <sub>2</sub> arrest	19.3	16.2	31.6	34.7 Cell Death		11.1	25.1	17.8	48.1 G <sub>2</sub> arrest		15.1	46.0	14.7	26.0 Cell Death	
MK-2	0	0.4	40.1	28.4	32.1	2.1	45.6	21.0	32.6		4.7	44.2	13.7	36.8		6.0	46.4	9.3	37.5	
	50	0.3	42.7	26.2	31.7	1.8	44.1	23.8	31.4		1.3	47.2	13.8	37.4		9.4	45.3	7.6	37.1	
	100	0.6	45.2	22.4	32.4	1.8	43.3	23.6	32.4		1.7	47.2	16.0	34.5		3.6	49.8	8.2	37.6	
	250	2.4	46.7	14.3	36.7	3.2	37.7	23.8	36.4 G <sub>2</sub> arrest		1.4	52.8	11.1	34.3		4.2	41.4	11.2	42.5 G <sub>2</sub> arrest	
	500	3.9	26.3	10.1	60.0 G <sub>2</sub> arrest	18.8	17.8	21.6	43.1 Cell death		2.5	39.4	11.3	46.5 G <sub>2</sub> arrest		5.2	44.5	15.4	33.6 S-phase	
MK-4	0	0.8	42.0	26.9	31.7	2.3	49.9	21.6	27.4		4.1	44.0	12.5	39.4		5.5	45.7	7.4	41.4	
	50	0.5	42.0	26.9	32.0	1.9	44.7	22.3	32.3		4.5	43.9	11.4	40.7		4.7	51.4	12.3	31.6	
	100	0.4	43.2	25.4	32.5	2.5	45.3	22.6	30.6		2.0	41.4	13.6	44.1		6.0	52.3	12.5	29.4	
	250	0.5	45.7	24.6	30.5	6.0	40.0	23.0	32.0		3.9	36.2	14.1	46.9		7.0	53.2	11.9	27.6	
	500	1.1	47.5	18.6	33.9 Slight $\Delta$	6.1	27.8	22.8	44.4 G <sub>2</sub> arrest		9.6	29.0	15.7	46.5 G <sub>2</sub> arrest		3.4	53.7	14.1	29.1 G <sub>1</sub> arrest	

cantharidin is greater than for its analogues. Interestingly, cantharidin also showed slight selectivity towards the colon cancer cells.

If the protein phosphatase inhibitor abrogates the G<sub>2</sub> checkpoint then the cells will not arrest in the G<sub>2</sub> phase of the cell cycle and the cells will continue through the cell cycle and accumulate in the G<sub>1</sub> phase of the cell cycle only. Similarly if the protein phosphatase inhibitors abrogates the G<sub>1</sub> checkpoint then the cells will not arrest in the G<sub>1</sub> phase of the cell cycle and accumulate in the G<sub>2</sub> phase of the cell cycle only. Cell cycle analysis using propidium iodide labelling of DNA has been used extensively in our laboratory to assess the effect of specific anticancer agents that induce S-phase cell cycle arrest and apoptotic cell death (Sakoff, Ackland and Stewart, 1998). Experiments were performed on a Becton Dickinson FACScan and using Cell Quest software.

Data shown in Table 3 and Figure 5 show the cell cycle response of L1210, HL60, HT29 and HCT116 cells. The cells were treated with 6Gy of radiation and then treated with cantharidin 6h later. The ability to abrogate cell cycle arrest was assessed 12h after the addition of the drugs. Cantharidin and MK-2 both abrogated radiation induced G<sub>1</sub> arrest in all cell lines. MK-4 also abrogated G<sub>1</sub> arrest in L1210, HL60 and HCT116 cells. In HT29 cells, MK-4 induced abrogation of the G<sub>2</sub> checkpoint. It is important to note that the exposure of HT29 cells to MK-4 induced the greatest cytotoxicity (IC<sub>50</sub> 14µM) as determined by the MTT assay. Not surprisingly, the ability to abrogate the G<sub>2</sub> checkpoint was more lethal than the ability to abrogate the G<sub>1</sub> checkpoint.

Table 3

## Checkpoint Abrogation

Cell Cycle Distribution (percentage of total) of tumour cell lines 18h after 6Gy of radiation and 12h after cantharidin or cantharidin analogue treatment.

Method : Flow Cytometry of Propidium Iodide stained cells.

Agent	L1210 cells					HL60 cells					HCT116 cells					HT29 cells				
	$\mu$ M	sub G <sub>1</sub>	G <sub>1</sub>	S	G <sub>2</sub> +M	sub G	G <sub>1</sub>	S	G <sub>2</sub> +M	sub G	G <sub>1</sub>	S	G <sub>2</sub> +M	sub G	G <sub>1</sub>	S	G <sub>2</sub> +M			
Cantharidin	0	1.6	25.3	35.8	38.8	6.6	5.3	3.2	85.3	4.9	26.8	8.7	60.1	5.9	40.7	9.2	44.7			
	1	1.6	27.2	25.6	37.5	6.2	5.2	3.0	85.8	4.2	26.1	13.8	58.2	16.6	35.2	10.6	38.2			
	5	2.4	25.8	31.9	41.5	4.4	5.7	3.5	86.8	4.0	23.6	10.4	63.3	5.3	38.6	10.6	46.3			
	10	3.4	24.9	29.4	43.7	5.3	5.6	4.3	85.1	4.2	25.7	9.4	62.2	6.4	21.1	12.3	60.8			
	50	4.9	4.1	15.6	77.4	G <sub>1</sub> abrogation	14.3	10.2	11.3	64.9	Cell Death	12.0	12.2	15.6	63.3	G <sub>1</sub> abrogation	14.7	23.0	20.7	43.1
MK-2	0	1.6	16.4	31.1	52.1	7.0	5.9	2.2	85.1	3.7	30.8	7.9	57.2	17.3	31.8	8.7	41.3			
	50	4.0	19.0	27.8	50.1	5.9	6.1	2.8	85.5	3.3	32.8	6.3	57.3	10.3	35.4	8.8	44.7			
	100	3.5	18.4	23.0	55.8	5.5	6.1	3.3	85.4	3.1	29.9	7.2	59.6	3.5	40.6	9.0	45.9			
	250	6.9	11.2	10.0	71.9	8.1	5.4	2.8	83.9	6.2	23.9	4.3	65.4	2.7	24.9	12.3	59.4			
	500	5.4	3.4	2.9	88.4	G <sub>1</sub> abrogation	11.8	4.4	4.1	80.0	Cell Death	6.4	15.4	4.9	73.0	G <sub>1</sub> abrogation	8.8	24.1	20.4	45.2
MK-4	0	1.9	20.2	29.7	50.0	8.7	5.7	2.0	83.9	10.3	31.4	6.2	52.1	7.0	35.1	9.3	48.4			
	50	1.8	21.2	28.5	50.3	8.9	6.2	2.8	82.3	6.3	26.7	5.8	61.3	6.8	28.9	16.4	48.4			
	100	2.4	22.0	27.4	49.7	9.8	6.2	3.4	80.8	3.3	18.4	9.7	69.3	6.3	33.3	17.2	43.3			
	250	3.1	21.2	24.6	52.7	9.3	5.8	3.1	82.2	8.2	16.3	8.2	67.8	10.3	35.2	17.3	36.5			
	500	5.0	18.2	16.0	61.8	G <sub>1</sub> abrogation	11.6	5.2	5.4	78.2	Cell Death	14.9	13.1	10.6	61.9	G <sub>1</sub> abrogation	3.9	39.4	19.2	37.7



- 30 -

## (c) Combination studies:

The cell lines listed above are exposed continuously to cisplatin and the phosphatase inhibitor in various drug ratio combinations for 72h and then assayed for cytotoxicity. Similarly, the cells are exposed to 8 Gy of radiation and incubated with the phosphatase inhibitor and assessed for cytotoxicity at 72 h.

Data shown in Figures 6-9 shows the results of combination studies utilising the Median Effect Method in HT29 and HCT116 human colon cells. This method tests the cytotoxicity of various drug combinations from which a combination index can be calculated. A value of greater than one indicated antagonism, a value equal to 1 indicates additivity, while a value less than one indicates synergism. The HT29 and HCT116 cell lines were chosen as they have differing p53 status and they represent the tumour types that responded the greatest to cantharidin and its analogues.





The data show that the simultaneous combination of cisplatin and MK-4 in both HCT116 and HT29 cells was additive and not synergistic using drug molar ratios of 1:1, 10:1 and 1:10. An additive response indicated that the drugs were mediating their effects via two separate biochemical pathways. The simultaneous combination of taxotere and MK-4 in HT29 cells was also additive using drug molar ratios of 1:10, 1:100, 1:1000 (Taxotere: MK-4). However, this drug combination of taxotere and MK-4 induced a synergistic response in HCT116 cells. A synergistic response indicates that the two drugs were interacting in such a way as to enhance the overall cytotoxic response and to induce "more than the additive" response of each individual agent. Consequently, the addition of subtoxic levels of MK-4 clearly enhanced the cytotoxicity of taxotere.






Example 8Results and Discussion

Anhydrides and simple analogues were synthesised according to literature procedures (Eggelte et. al; 1973), and then subjected to a PP1 and PP2A bio-assay (see

5 biochemistry) to determine their ability to inhibit these enzymes. The results of initial screening at 100 mM are shown in Table 4, along with IC<sub>50</sub> values in some instances.

Table 4 The inhibition of protein phosphatase 1 and 2A by anhydride modified cantharidin analogues.

Compound	Inhibition of PP1 (%)	Inhibition of PP2A (%)	Selectivity PP2A/PP1
 1	90 IC <sub>50</sub> 2.4 μM	97 IC <sub>50</sub> 2.1 μM	0.875
 2	ND	95	
 3	46 IC <sub>50</sub> 50μM	6 IC <sub>50</sub> >10,000μM	>200
 4	13	11	

 5	15	8	
 6	9	11	
 7	ND	21	
 8	ND	15	
 9	ND	4	





Of the compounds listed in Table 4, only 1 and 2 show any significant inhibition of PP2A. at 97% and 95% respectively (with little selectivity apparent for either enzyme). Interestingly the bioisosteric replacement of the anhydride oxygen atom of 1 results in a complete loss of inhibition. Indeed no modification of the cyclic anhydride, is tolerated, and consequently results in no inhibition of PP2A.



Previously we have shown that analog 2 undergoes a rapid conversion to the dicarboxylic acid under assay conditions. We thus examined the stability of the non-

active analogues (in Table 4) and found that they were stable under assay conditions showing no decomposition, in fact 5 can be synthesised via the Diels-Alder reaction in water (Eggelte et al; 1973).


In all instances, the corresponding dicarboxylic acid derivatives display lower inhibitory values at PP2A (Tables 5 and 6). Even though the anhydrides undergo a facile ring opening to the dicarboxylic acids, the original conformation presented at the active site must also play a role in determining the overall level of inhibition. Consequently, we believe that the conformation of anhydride carbonyl groups is more favourable for inhibition (essentially only one conformation presented at the active site), than that of the dicarboxylic acid (four possible minimum energy conformations, data not shown).

Table 5 Effects of anhydride to dicarboxylic acid on the inhibition of PP2A

Entry	Anhydride	Inhibition (%)	Carboxylic acid	Inhibition (%)
1		97 (This work)		80
2		92-95		92-95





3		48		17
---	---	----	--	----

In an attempt to determine the feasibility of anhydride opening via nucleophilic attack from Tyr272, we conducted a series of model experiments in which 2 was allowed to stand in a chloroform solution of phenol. This mixture was examined periodically by <sup>1</sup>H NMR spectroscopy and showed the growth of a new species over a period of time (ca 10 days). Further analysis indicated the presence of a phenolate ester of norcantharidin (scheme 4). Consequently, a metal assisted or nucleophilic attack under physiological conditions represents a possible mode of assisted ring opening with the anhydride held in a favourable conformation within the active site. In turn the resultant diacid rapidly binds in a more favourable manner.



Scheme 4

Table 6 Inhibition of PP1 and PP2A by selected cantharidin analogues

Entry	Compound	Inhibition of PP1 (%)	Inhibition of PP2A (%)	Selectivity PP2A/PP1
1		90 (IC <sub>50</sub> 2.4μM)	97 (IC <sub>50</sub> 2.1μM)	0.875
2		46 (IC <sub>50</sub> 50μM)	6 (IC <sub>50</sub> >10000μM)	>200
4		3	3	Not determined
5		15	69	Not determined

The results presented herein indicate that cantharidin analogues, via anhydride opening are more potent inhibitors of PP2A. Analogues in which the anhydride moiety has been modified preventing a facile ring opening (except where otherwise indicated) are extremely poor inhibitors of PP2A (Tables 5 and 6).

However, the most interesting result reported herein (see table 4) is the selective inhibition of PP1 by the dimethyl ester (3). Simple diesterification of 2 has completely reversed the previously reported PP2A selectivity (ca 10 fold) of norcantharidin for PP2A to yield selective small synthetic molecule for the inhibition of either PP1 or

PP2A. Again this suggests that presentation of a diacid moiety to the active site is crucial for the inhibition of PP2A. No such restrictions are apparent with the limited structure activity data for PP1.

A synthetic inhibitor such as 3 represents a significant advance on the currently  
5 widespread inhibitors of PP1 and PP2A.

In conclusion, the present inventors have demonstrated that a facile ring opening of the anhydride moiety is relevant for inhibition at PP2A. Also, that modification of the dicarboxylic acid moiety gives rise to a PP1 selective compound.

The above describes some embodiments of the present invention. Modifications  
10 obvious to those skilled in the art can be made without departing from the scope of this invention.

#### Industrial Applicability

It should be clear that the present invention will find light applicability, especially in the medical and veterinary fields.

## REFERENCES

- Walsh A H, Chen A T, Honaken R E, *FEBS Lett.*, 1997, 416, 230
- Roberge M, Tudan C. Hung SMF, Harder KW, Jirik FR. and Anderson H *Cancer Research* 1994, 54, 6115-6121.
- 5 McCluskey, A; Taylor, C; Quinn, R J; Suganuma, M; Fujiki, H; *Bioorg. Med. Chem. Lett.*, 1996, 6, 1025.
- McCluskey A, Keane M A, Sim A T R, *J. Med. Chem.* 1998, submitted.
- Enz, A; Zenke, G; Pombo-Villar; *Bioorg. Med. Chem. Lett.*, 1997, 7, 2513; and
- Sodeoka, M; Baba, Y; Kobayashi, S; Hirukawa, N; *Bioorg. Med. Chem. Lett.*, 1997, 7,
- 10 1833.
- Cohen, P; Cohen, PTW; *J. Biol. Chem.*, 1989, 264, 21435.
- Cohen, P; "The Structure and Regulation of Protein Phosphatases" Raven Press: New York, 1990, p230-235.
- Shenolinkar, S; Mairn, A C; *Adv. Second Messenger Phosphoprotein Res.*, 1991b, 23, 1.
- 15 Krebs, E G; *Angew. Chem., Int. Ed. Engl.*, 1993, 32, 1122.
- Fischer, E; *Angew. Chem., Int. Ed. Engl.*, 1992, 32, 1130.
- Tachibana, K; Scheuer, P J; Tsukitani, Y; Kikuchi, H; Van Engen, D; Clardy, J;
- Gopichand, Y; Schmitz, F J; *J. Am. Chem. Soc.*, 1981, 103, 2469.
- Cheng, X-C; Kihara, T; Kusakabe, H; Magae, J; Kobayashi, Y; Fang, R-P; Ni, Z-F;
- 20 Shen, Y-C; Ko, K; Yamaguchi, I; Isono, K; *J. Antibiot.*, 1987, 40, 907.
- Botes, D P; Tuinman, A A; Wessels, P L; Viljoen, C C; Kruger, H; Williams, D H;
- Santikarn, S; Smith, K J; Hammond, S J; *J. Chem. Soc. Perkin Trans I*, 1984, 2311.



- Herada, K-I; Matsuura, K; Suzuki, M; Oka, H; Watanabe, M F; Oishi, S; Dahlem, A M;  
Beasley, V R; Carmichael, W W; *J. Chromatog.*, 1998, 448, 275.
- Honaken, R E; *FEBS Lett.*, 1993, 330, 283.
- Matszawa, S; Suzuki, T; Suzuki, M; Matsuda, A; *FEBS Lett.*, 1994, 356, 272.
- 5 Wang, G-S; *J. Ethnopharmacol.*, 1989, 26, 147.
- Cavill, G W K; Clark, D V; *Naturally Occuring Insecticides* (M. Jacobson and D G  
Crosby, EDS), Marcel Dekker, New York, 1971, 271.
- Oaks, W W; DiTunno, J F; Magnani, T; Levey, H A; Mills, L C; *A.M.A. Arch. Int.  
Med.*, 1960, 105, 106.
- 10 Goldfarb, M T; Gupta, A K; Sawchuk, W S; *Dermatologic Clinics*, 1991, 9, 287.
- A. Schmitz, D G; *J. Vet. Intern. Med.*, 1989, 3, 208.
- Polettini, A; Crippa, U; Ravagii, A; Saragoni, A.; *Forensic. Sci. Int.*, 1992, 56(1), 37.
- Li, Y-M; Casida, J E; *Proc. Natl. Acad. Sci. USA*, 1992, 89, 11867.
- McCluskey, A; Taylor, C; Quinn, R J; Suganuma, M; Fujiki, H; *Bioorg. Med. Chem.*  
15 *Lett.*, 1996, 6, 1025.
- Eggelte, T A; de Koning, H; Huisman, H O; *Tetrahedron*, 1973, 29, 2445; 3 see:  
Tamura, Y; Imanishi, K; Miki, Y; Ikeda, M; *Synthesis*, 1997, 559; 5 & 9 see: Kwart, H;  
Burchuk, I; *J. Am. Chem. Soc.*, 1952, 74, 3094.
- Goldberg, J; Huang, H-B;; Kwon, Y-G; Greengard, P; Nairn, A C; Kuriayan, J; *Nature*,  
20 1995, 376, 745.
- Quinn, R J; Volter, K; Embury, K J, *unpublished results*.

- Cohen, P; Alemany, S.; Hemmings, B A; Resink, T J; Stralfors, P and Tung, H Y L in *Methods in Enzymology* (Ed. J D Corbin and R A Johnson) 1988, 159, 390, Academic Press (London).
- Hardie, DG; Haystead TAJ; Sim ATR; Protein Phosphorylation, in methods in  
 5 enzymology, in press. *Enz. A*; Zenke, G; Pombo-Villar; *Bioorg. Med. Chem. Lett.*, 1997, 7, 2513.
- Sodeoka, M; Baba, Y; Kobayashi, S; Hirukawa, N; *Bioorg. Med. Chem. Lett.*, 1997, 7, 1833.
- Durfee T., et al., (1993). The retinoblastoma protein associates with the protein  
 10 phosphatase type 1 catalytic subunit. *Genes Dev.* 7, 555-569.
- Ghosh, S. et al. (1996). Okadaic acid overrides the S-phase checkpoint and accelerates progression of G2-phase to induce premature mitosis in Hela cells. *Exp. Cell Res.* 227:165-169.
- Herwig, S., and Strauss, M. (1997). The retinoblastoma protein: a master regulator of  
 15 cell cycle, differentiation and apoptosis. *Eur.J.Biochem.* 246:581-601.
- Jackman, AL and Calvert. AH (1995). Folate-based thymidylate synthase inhibitors as anticancer drugs. *Ann. Oncol.* 6:871-881, 1995.
- Kawamura K-I, Grabowski D, Weizer K, Bukowski R, and Ganapathi R (1996).  
 Modulation of vinblastine cytotoxicity by dilantin (phenytoin) or the protein phosphatase  
 20 inhibitor okadaic acid involves the potentiation of anti-mitotic effects and induction of apoptosis in human tumour cells. *Br. J. Cancer* 73: 183-188.
- Lazzereschi, D. et al. (1997). The phosphatase inhibitor okadaic acid stimulates the TSH-induced G1-S phase transition in thyroid cells. *Exp. Cell Res.* 234:425-433.

- Nakamura K. and Antoku S. (1994). Enhancement of X-ray cell killing in cultured mammalian cells by the protein phosphatase inhibitor calyculin A. *Cancer Res* 54:2088-20990.
- O'Connor PM (1996). Cell cycle checkpoints: Targets for anti-cancer therapy. *Anti-Cancer*  
5 *Drugs* 7(sup 3): 135-41.
- O'Connor PM (1997). Mammalian G1 and G2 phase checkpoints. *Cancer Surveys* 29:151-182.
- Peters GJ and Ackland SP (1996). New antimetabolites in preclinical and clinical development. *Exp. Opin. Invest. Drugs* 5, 637-679.
- 10 Schšnthal A (1992). Okadaic acid - a valuable new tool for the study of signal transduction and cell cycle regulation? *New Biol.* 4, 16.
- Stein, G.S. et al. (1998). The molecular basis of cell cycle and growth control. Wiley-Liss Publishers. New York.
- Yamashita K., et al (1990). Okadaic acid, a potent inhibitor of type 1 and type 2A protein  
15 phosphatases, activates cdc2/H1 kinase and transiently induces a premature mitosis-like state in BHK21 cells. *EMBO J.* 9: 4331-4338.

09/743841

09/743841


JC07 Rec'd PCT/PTO 16 JAN 2001

U.S. CLAIMS


- 41 -

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:-

1. A cell permeable inhibitor of protein phosphatase, said inhibitor being an anhydride modified cantharidin analogue.
2. An inhibitor according to claim 1, wherein the phosphatase is phosphatase 1 and/or phosphatase 2A.
3. An inhibitor according to claim 1 or 2 wherein the anhydride modified cantharidin analogue is oxidatively stable.
4. A compound of the formula:



- 10 wherein  $R_1$  and  $R_2$  are H, aryl or alkyl; X is O, N or S; Y is O, S, SR, NH, NR,  $CH_2OH$ ,  $CH_2OR$ ; R is alkyl or aryl; A and B are H or  $CH_3$ ; W and Z are  $CHOH$  or  $C=O$  and  $R_1$  and  $R_2$  can cyclise to form a ring as follows:



wherein  $R_3$  and  $R_4$  are H, aryl or alkyl.

- 15 5. A compound according to claim 3, wherein the aryl group is phenyl or naphthyl and wherein the aryl group is attached via a carbon spacer of between 6 and 10 carbon atoms.
6. A compound according to claim 3 or claim 4, wherein the alkyl group is  $C_1$ - $C_{10}$ .

- 42 -

7. A process for producing anhydride modified cantharidin analogues for use in the treatment of cancer or for the sensitising cancer cells to one or more cancer treatments comprising the step of reacting a diene with an ene.

8. A process according to claim 7 further comprising hydrogenation of the adduct  
5 of the diene and the ene.

9. A process according to claim 7 further comprising ring opening of the adduct of the diene and the ene.

10. A process for producing anhydride modified cantharidin analogues, said process including the steps of:

10 dissolving a diene in a suitable solvent and adding to the resultant solution an ene.

11. A process for producing anhydride modified cantharidin analogue, said process including the steps of:

15 dissolving a furan in a suitable solvent and adding to the resultant solution an ene;

incubating the solution at a temperature and for a time sufficient to form a precipitate; and

collecting the precipitate and recrystallising the analogue.

12. A process for producing anhydride modified cantharidin analogue, said process  
20 including the steps of:

mixing thiophene and maleic anhydride at room temperature in a suitable solvent;

00507434105004

- 43 -

compressing the mixture at a temperature and pressure sufficient to facilitate a reaction to take place; and

purifying the analogue.

13. A method of treating cancer which method comprises administering to a patient  
5 in need of such treatment, an effective amount of an inhibitor according to claim 1 or a compound according to claim 4, together with a pharmaceutically acceptable carrier, diluent and/or excipient.

14. A method according to claim 13, wherein the cancer is inherently resistant to conventional chemotherapy.

10 15. A method according to claim 13, wherein the cancer is colon cancer or non small-cell lung cancer.

16. A method according to claim 13, wherein the inhibitor or the compound is administered intravenously.

17. A method of sensitising cancer cells to at least one method of treating cancer,  
15 which method of sensitising comprises administering to a patient in need of such treatment, an effective amount of an inhibitor according to claim 1 or a compound according to claim 4, together with a pharmaceutically acceptable carrier, diluent and/or excipient.

18. A method according to claim 17, wherein the at least one cancer treatment is  
20 selected from treatments involving irradiation and anti-cancer agents.

19. A method according to claim 17, wherein the cells have deficient p53 activity.

09743841-050001

- 44 -

20. A method of treating cancer which method comprises:

administering to a patient in need of such treatment, an effective amount of an anhydride modified cantharidin analogue of claim 1 or a compound according to claim 4 to sensitise cells of the patient to one or more cancer treatments; and

5 utilising the one or more cancer treatments.

21. A method of screening compounds for use in sensitising cancer cells to at least one method of treating cancer, and comprising:

screening for anti-cancer activity; and

10 screening for ability to abrogate either the G<sub>1</sub> or the G<sub>2</sub> checkpoint of the cancer cell cycle.

22. A method according to claim 21 further comprising the step of screening for the ability of the compounds of sensitise cancer cells to one or more cancer treatments.

23. A method according to claim 21 wherein the one or more cancer treatments are selected from treatments involving cisplatin, irradiation, taxanes and antimetabolites.

15 24. A method according to claim 21 wherein the screening is conducted on haematopoietic cells or solid tumour cells, having varying p53 activity.

25. A method according to claim 24, wherein the cells are selected from the group consisting of L1210 (murine leukaemia, p53 wildtype), HL60 (human leukaemia, p53 nul), A2780 (human ovarian carcinoma, p53 wildtype), ADDP (cisplatin resistant


20 A2780 cells, p53 mutant), SW480 (human colon carcinoma, p53 mutant), WiDr (human colon carcinoma, p53 mutant), HT29 (human colon carcinoma, p53 mutant), HCT116 (human colon carcinoma, p53 wildtype) and 143B (human osteosarcoma, p53 mutant).

0943841-050904

- 45 -

26. A compound selected from a group comprising compounds (a) to (n) below:


(a)




(b)




(c)




(d)



(e)




(f)




(g)




(h)



(i)



(j)




09743344.050504




- 46 -


(k)




(l)



(m)



(n)



27. Use of an inhibitor according to claim 1, or a compound according to claim 4 for the manufacture of a medicament for the treatment of cancer.

28. Use according to claim 27, wherein the cancer is colon cancer or non small-cell lung cancer.

29. Use according to claim 28, wherein the medicament is administered intravenously.

0574344-050301

- 1/27 -




Figure 1

# Cantharidin and Cantharidin Analogues

WO 00/04023

09/743843

PCT/AU99/00567

- 2/27 -




Figure 2

MK-7, MK-8, MK-9: isomers of MK-2

- 3/27 -




Figure 3

- 4/27 -




Figure 3 (cont'd)

- 5/27 -




Figure 3 (cont'd)

- 6/27 -




Figure 3 (cont'd)

- 7/27 -




Figure 3 (cont'd)



- 8/27 -




Figure 3 (cont'd)

- 9/27 -




Figure 3 (cont'd)

- 10/27 -




Figure 3 (cont'd)

- 11/27 -




Figure 3 (cont'd)

- 12/27 -




Figure 3 (cont'd)

- 13/27 -




Figure 3 (cont'd)

- 14/27 -





Figure 3 (cont'd)




Cell Cycle Analysis 12h following exposure to cantharidin, MK-2 or MK-4

Figure 4



- 16/27 -



Cell Cycle Analysis 18h after 6Gy of radiation and 12h after exposure to cantharidin, MK-2 or MK-4

Figure 5



**Combination Index vs Fn Affected: HCT-116 cells, simultaneous x72hr, CDDP & MK-4 various ratios, MTT**





Figure 6(b)

**Mhc99423.xls**




- 20/27 -



Mh990416.xls

Figure 7(a)


# Combination Index vs Fn Affected: HT29cells, simultaneous x72hr, CDDP & MK-4 various ratios, MTT



Mh990423.xls

Figure 7(b)


- 22/27 -



Mhc99514.xls

Figure 8(a)

**Combination Index vs Fn Affected: HCT-116cells,  
simultaneous x72hr, TXT & MK-4 various ratios, MTT**




**Mhc99521.xls**

Figure 8(b)



- 24/27 -


**Combination Index vs Fn Affected: HCT-116cells,  
simultaneous x72hr, TXT & MK-4 various ratios, MTT**



**Mhc99528.xls**

Figure 8(c)


- 25/27 -



Mht99514.xls

Figure 9(a)


# Combination Index vs Fn Affected: HT29 cells, simultaneous x72hr, TXT & MK-4 various ratios, MTT



Mht99521.xls

Figure 9(b)

- 27/27 -



Mht99528.xls

Figure 9(c)

As a below named inventor, I hereby declare that my residence, post office address and citizenship are as stated below next to my name, and I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the **INVENTION ENTITLED ANHYDRIDE MODIFIED CANTHARIDIN ANALOGUES USEFUL IN THE TREATMENT OF CANCER**

the specification of which (CHECK applicable BOX(ES))

X A. ☐ is attached hereto.  
BOX(ES) → B. ☒ was filed on 16 January 2001 as U.S. Application No. 09/743,841  
→ C. ☒ was filed as PCT International Application No. PCT/ AU99/00567 on 14 July 1999

and (if applicable to U.S. or PCT application) was amended on

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above. I acknowledge the duty to disclose all information known to me to be material to patentability as defined in 37 C.F.R. 1.56. Except as noted below, I hereby claim foreign priority benefits under 35 U.S.C. 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT International Application which designated at least one other country than the United States, listed below and have also identified below any foreign application for patent or inventor's certificate, or PCT International Application, filed by me or my assignee disclosing the subject matter claimed in this application and having a filing date (1) before that of the application on which priority is claimed, or (2) if no priority claimed, before the filing date of this application:

**PRIOR FOREIGN APPLICATION(S)**

Number	Country	Day/MONTH/Year Filed	Date first Laid-open or Published	Date Patented or Granted	Priority NOT Claimed
PP 4665	Australia	14 July 1998			

I have prior foreign applications, X box at bottom and continue on attached page.

Except as noted below, I hereby claim domestic priority benefit under 35 U.S.C. 119(e) or 120 and/or 365(c) of the indicated United States applications listed below and PCT international applications listed above or below and, if this is a continuation-in-part (CIP) application, insofar as the subject matter disclosed and claimed in this application is in addition to that disclosed in such prior applications, I acknowledge the duty to disclose all information known to me to be material to patentability as defined in 37 C.F.R. 1.56 which became available between the filing date of each such prior application and the national or PCT international filing date of this application:

**PRIOR U.S. PROVISIONAL, NONPROVISIONAL AND/OR PCT APPLICATION(S)**

Application No. (series code/serial no.)	Day/MONTH/Year Filed	Status	Priority NOT Claimed
		pending, abandoned, patented	

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

And I hereby appoint Pillsbury Winthrop LLP, Intellectual Property Group, 1100 New York Avenue, N.W., Ninth Floor, East Tower, Washington, D.C. 20005-3918, telephone number (202) 861-3000 (to whom all communications are to be directed), and the below-named persons (of the same address) individually and collectively my attorneys to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith and with the resulting patent, and I hereby authorize them to delete names/numbers below of persons no longer with their firm and to act and rely on instructions from and communicate directly with the person/assignee/attorney/firm/ organization who/which first sends/sent this case to them and by whom/which I hereby declare that I have consented after full disclosure to be represented unless/until I instruct the above Firm and/or a below attorney in writing to the contrary.

Paul N. Kokulis	16773	Kendrew H. Colton	30368	Roger R. Wise	31204	Anthony L. Miele	34393
G. Lloyd Knight	17698	G. Paul Edgell	24238	Michael R. Dzwonczyk	36787	Robert J. Walters	40862
Kevin E. Joyce	20508	Lynn E. Eccleston	35861	W. Patrick Bengtsson	32456	Brian J. Beatus	38825
George M. Sirilla	18221	Timothy J. Klima	34852	Jack S. Barufka	37087	John R. Wetherell	31678
Donald J. Bird	25323	David A. Jakopin	32995	Adam R. Hess	44835	Alan E. Dow	35123
Dale S. Lazar	28872	Mark G. Paulson	30793	William P. Atkins	38821	Jane K. Babin	47224
Paul E. White, Jr.	32011	Stephen C. Glazier	34361	Paul L. Sharer	36004	Robert M. Bedgood	43488
Glenn J. Perry	28458	Richard H. Zaitlen	27248	Robin L. Teskin	35030		

**(1) INVENTOR'S SIGNATURE:**

Date: 10<sup>th</sup> March 2001

Adam	McCluskey		
First	Middle Initial	Family Name	
Residence	Mayfield	Australia	Great Britain
City	State/Foreign Country	Country of Citizenship	
Mailing Address	19 Simpson Court		
(include Zip Code)	NSW 2304		

**(2) INVENTOR'S SIGNATURE:**

Date: 19<sup>th</sup> March 2001

Jennette	A	Sakoff	
First	Middle Initial	Family Name	
Residence	Belmont	Australia	Australia
City	State/Foreign Country	Country of Citizenship	
Mailing Address	64 Gilsmore Street	71 Green Point Drive	
(include Zip Code)	NSW 2280		

"X" box ☒ FOR ADDITIONAL INVENTORS, and proceed on the attached page to list each additional inventor.  
☐ See additional foreign priorities on attached page (incorporated herein by reference).

Atty. Dkt. No. 80905-272441  
(M#)

## DECLARATION AND POWER OF ATTORNEY

(continued)

## ADDITIONAL INVENTORS:

MAY 09 2001

Date:

19/3/01

## (3) INVENTOR'S SIGNATURE:

Stephan		Ackland	
First		Middle Initial	
Residence		Family Name	
New Lambton Heights		Australia	
City		State/Foreign Country	
Mailing Address		Country of Citizenship	
(include Zip Code)		95 Carrington Parade	
		NSW 2305	

## (4) INVENTOR'S SIGNATURE:

Date:

22/3/01

Alistair		T.R. Sim	
First		Middle Initial	
Residence		Family Name	
Callaghan		Australia	
City		State/Foreign Country	
Mailing Address		Country of Citizenship	
(include Zip Code)		The University of Newcastle, The Faculty of Medical and Health Sciences, Dept. of Medical Biochemistry, University Drive	
		NSW 2308	

## (5) INVENTOR'S SIGNATURE:

Date:

First		Middle Initial	
Residence		Family Name	
City		State/Foreign Country	
Mailing Address		Country of Citizenship	
(include Zip Code)			

## (6) INVENTOR'S SIGNATURE:

Date:

First		Middle Initial	
Residence		Family Name	
City		State/Foreign Country	
Mailing Address		Country of Citizenship	
(include Zip Code)			

## (7) INVENTOR'S SIGNATURE:

Date:

First		Middle Initial	
Residence		Family Name	
City		State/Foreign Country	
Mailing Address		Country of Citizenship	
(include Zip Code)			

## (8) INVENTOR'S SIGNATURE:

Date:

First		Middle Initial	
Residence		Family Name	
City		State/Foreign Country	
Mailing Address		Country of Citizenship	
(include Zip Code)			

## (9) INVENTOR'S SIGNATURE:

Date:

First		Middle Initial	
Residence		Family Name	
City		State/Foreign Country	
Mailing Address		Country of Citizenship	
(include Zip Code)			